



## King's Research Portal

DOI:

[10.1016/j.msard.2017.12.011](https://doi.org/10.1016/j.msard.2017.12.011)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Rhone, E. E., Cho, P. S. P., Birring, S. S., Galloway, J., & Silber, E. (2017). Pulmonary Sarcoidosis in a patient with Multiple Sclerosis on Daclizumab monotherapy. *Multiple Sclerosis and Related Disorders*, 20, 25-27. <https://doi.org/10.1016/j.msard.2017.12.011>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.


- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

Pulmonary Sarcoidosis in a patient with Multiple Sclerosis on Daclizumab monotherapy

Elijah Edward Rhone, Peter Siu Pan Cho, Surinder S. Birring, James Galloway, Eli Silber



PII: S2211-0348(17)30349-8  
DOI: <https://doi.org/10.1016/j.msard.2017.12.011>  
Reference: MSARD725

To appear in: *Multiple Sclerosis and Related Disorders*

Received date: 9 November 2017  
Revised date: 19 November 2017  
Accepted date: 15 December 2017

Cite this article as: Elijah Edward Rhone, Peter Siu Pan Cho, Surinder S. Birring, James Galloway and Eli Silber, Pulmonary Sarcoidosis in a patient with Multiple Sclerosis on Daclizumab monotherapy, *Multiple Sclerosis and Related Disorders*, <https://doi.org/10.1016/j.msard.2017.12.011>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## **Pulmonary Sarcoidosis in a patient with Multiple Sclerosis on Daclizumab monotherapy**

Authors: Elijah Edward Rhone<sup>a</sup>; Peter Siu Pan Cho<sup>b</sup>; Surinder S. Birring<sup>bc</sup>; James Galloway<sup>c</sup>; Eli Silber<sup>a</sup>

<sup>a</sup>Department of Neurology, King's College Hospital, Denmark Hill, London, SE5 9RS, UK

<sup>b</sup>Division of Asthma, Allergy and Lung Biology, King's College London, Weston Education Centre, Denmark Hill, London SE5 9RJ, UK

<sup>c</sup>Department of Respiratory Medicine, King's College Hospital, Denmark Hill, London, SE5 9RS, UK

<sup>d</sup>Department of Rheumatology, King's College London King's College London, Weston Education Centre, Denmark Hill, London SE5 9RJ, UK

<sup>e</sup>Department of Rheumatology, King's College Hospital, Denmark Hill, London SE5 9RS, UK

Corresponding author:

Dr Elijah Rhone

Clinical Research Fellow in Neurology

King's College Hospital,

Denmark Hill,

London, England

SE5 9RS

Phone: (+44) 2032991905

Fax: (+44) 2032993587

Email: [erhone@nhs.net](mailto:erhone@nhs.net)

Co-author contact details

Peter Cho: [p.cho@nhs.net](mailto:p.cho@nhs.net)

Surinder Birring: [surinder.birring@nhs.net](mailto:surinder.birring@nhs.net)

James Galloway: [james.galloway@kcl.ac.uk](mailto:james.galloway@kcl.ac.uk)

Eli Silber: [eli.silber@nhs.net](mailto:eli.silber@nhs.net)

Word Count – 948 (including abstract but excluding figure description and references)

Number of references: 10

Number of figures: 2

Search Terms: Multiple sclerosis, Daclizumab, Sarcoidosis

Author contributions:

All authors were involved in the clinical care of this patient. Elijah Rhone completed the first draft of the manuscript. All authors were involved in critical revision of the manuscript.

# **Pulmonary Sarcoidosis in a patient with Multiple Sclerosis on Daclizumab monotherapy**

## **ABSTRACT**

As new immunomodulatory therapies continue to be licensed for use in Multiple Sclerosis, it is important to remain vigilant for new, unexpected associations relating to these medications. We highlight this by reporting on a case of a 45-year-old man who developed systemic, non-specific symptoms following long term use of daclizumab and was subsequently diagnosed with sarcoidosis. We go on to briefly discuss the action of daclizumab, in particular the effect it has on CD56<sup>bright</sup> natural killer cells, a cell type that has been investigated in relation to sarcoidosis.

## CASE DETAILS

A 45-year old Caucasian male presented with a 2-month history of fevers, night sweats and non-productive cough. He had been on a series of consecutive trials of daclizumab for Relapsing Remitting Multiple Sclerosis (RRMS), totalling 79 months (12-months placebo-controlled, 12-months dose-blinded, 55-months open-label of 150mg monthly subcutaneously). He had experienced three relapses in the three years prior to commencing the trials but remained relapse-free during them. He had no significant co-morbidity. Physical examination revealed new bilateral inguinal lymphadenopathy. Neurological examination demonstrated longstanding altered left-sided facial sensation, mild instability on Romberg testing; Expanded Disability Status Scale (EDSS) score was stable at 2.0. Investigations revealed elevated ESR (35mm/hr), ACE (62 IU/L), IgG (27.65 g/L), IgM (5.51 g/L) and total protein (91 g/L) but were otherwise unremarkable (lymphocyte count  $1.72 \times 10^9$  g/L, corrected-calcium 2.35 mmol/L). Daclizumab was suspended pending further investigation. CT-PET confirmed high avidity lymphadenopathy in the mediastinum and abdominal chains, but no visceral uptake (figure a). Lymph node aspirate obtained via endobronchial ultrasound demonstrated non-necrotising granulomatous inflammation (figure b).

Brain MRI before/after gadolinium was stable with no new or enhancing lesions and no meningeal enhancement. Two months following daclizumab cessation, spirometry demonstrated moderate obstruction; FEV1 2.13 (62.3% predicted), FVC 3.62 (83.7% predicted), FEV1/FVC ratio 58.8%. At this point, all symptoms except the non-productive cough had resolved. No alternative aetiology was identified and a clinical diagnosis of sarcoidosis was made. Inhaled corticosteroids were commenced followed by a short trial of oral corticosteroids, both of which had no effect on the cough.

CT-PET 6-months following daclizumab cessation demonstrated reduction in lymph node uptake. Repeat spirometry 10-weeks following discontinuation of corticosteroids was normal; FEV1 3.01 (88% predicted), FVC 3.74 (86.5% predicted), FEV1/FVC ratio 80.5%. Considering subjective and objective evidence of improvement following daclizumab cessation, it was reasoned that daclizumab was the probable cause and it was therefore permanently discontinued.

## DISCUSSION

Sarcoidosis is a multi-system granulomatous disorder that most commonly affects the pulmonary and lymphatic systems. In most patients, granulomas resolve spontaneously or following appropriate therapy (e.g. corticosteroids with or without steroid sparing agents such as methotrexate). A minority of patients develop chronic progressive disease, characterised by persisting granulomas, which may undergo fibrosis, causing irreversible organ damage. Sarcoid granulomas are characterised by monocytes, macrophages and activated T cells, with interleukin-2 (IL-2) a driver cytokine of this Th1-type cellular response.<sup>1</sup>

Daclizumab is a monoclonal antibody approved for treatment of RRMS. Trials have demonstrated its efficacy in reducing relapses, disability progression and development of new MRI lesions<sup>2</sup>. Daclizumab binds to the alpha chain (CD25) on high-affinity IL-2 receptors, inhibiting the association of IL-2 with the receptor. The IL-2 pathway is involved in activation, proliferation and differentiation of T cells and daclizumab is hypothesised to suppress IL-2-mediated T-cell expansion and activation<sup>2</sup>. The current understanding of sarcoid immunopathology makes its occurrence paradoxical in the setting of Daclizumab therapy. One speculative hypothesis lies in the effect of daclizumab on natural killer cells. CD56<sup>bright</sup> cells are a subset of natural killer (NK) cells, observed in small numbers in

peripheral blood but as a much greater proportion within lymph nodes<sup>3</sup>. These cells are immunoregulatory and have the capacity to produce a range of inflammatory cytokines (e.g. TNF- $\alpha$ , IFN- $\gamma$ ) when activated<sup>3</sup>. CD56<sup>bright</sup> NK cell populations expand significantly during daclizumab therapy. The proposed mechanism is that blockade of high-affinity IL-2 receptors leads to reduced IL-2 consumption as well as increased IL-2 production due to lack of negative feedback resulting in increased local availability of IL-2. CD56<sup>bright</sup> NK cells express an intermediate-affinity IL-2 receptor. Daclizumab does not interact with this receptor allowing IL-2 to bind freely, resulting in cell proliferation<sup>2</sup>.

Interestingly, higher numbers of CD56<sup>bright</sup> cells have been observed in broncho-alveolar lavage fluid of individuals with sarcoidosis compared to controls<sup>4</sup> and higher proportions of NK cells in this fluid have been found to correlate with a poor prognosis<sup>5</sup>. However, an expansion of CD56<sup>bright</sup> cells can be seen in a range of diseases and it is unclear if they are a driver of disease or a consequence of the pathological process<sup>3</sup>.

In a review of daclizumab, the Center for Drug Evaluation and Research (a division of the U.S. Food and Drug Administration), with access to raw trial data, identified nine cases of sarcoidosis amongst various RRMS trial cohorts, with another four potential cases. Based on these nine confirmed cases, the incidence of sarcoidosis was estimated to be approximately 1.32/1,000 patient years compared to a reported rate in the general population of 1.0-35.5 per 100,000/year<sup>6</sup>. One case of sarcoidosis has also been reported in a patient treated off study with another form of daclizumab in adult T-cell leukaemia/lymphoma<sup>7</sup>.

The development of sarcoidosis has also been observed following the use of other disease modifying therapies. Several cases have been reported following the use of interferon- $\beta$ ,

including specifically in patients being treated for multiple sclerosis<sup>8</sup>. There have also been reports of cases following the use of alemtuzumab<sup>9</sup> and natalizumab<sup>10</sup> in the treatment of other diseases; although to our knowledge, there are no reports in the literature of the development of sarcoidosis in patients treated with these medications for multiple sclerosis. This case emphasises the need to be vigilant for unexpected associations when using novel immunomodulatory therapies.



Figure a - PET- CT demonstrating metabolically active mediastinal lymphadenopathy






Figure b - Cytology photomicrograph demonstrating granulomas following EBUS sampling

### **Acknowledgements**

We would like to thank Dr Emma Mclean, consultant histopathologist at Guy's and St Thomas' Hospital for providing the cytology image.

### **Consent**

The patient provided written consent for the details of this case to be published in an anonymised fashion.

## References

1. Broos CE, van Nimwegen M, Hoogsteden HC, Hendriks RW, Kool M, Van den Blink B. Granuloma formation in pulmonary sarcoidosis. *Front Immunol*. 2013 Dec 10;4:437.
2. Wiendl H, Gross CC. Modulation of IL-2R $\alpha$  with daclizumab for treatment of multiple sclerosis. *Nat Rev Neurol*. 2013;9:394–404.
3. Poli A, Michel T, Thérésine M, Andrès E, Hentges F, Zimmer J. CD56bright natural killer (NK) cells: an important NK cell subset. *Immunology*. 2009 Apr;126(4):458–465.
4. Katchar K, Soderstrom K, Wahlstrom J, Eklund A, Grunewald J. Characterisation of natural killer cells and CD56+ T-cells in sarcoidosis patients. *Eur Respir J*. 2005 26(1):77–85.
5. Tutor-Ureta P, Citores MJ, Castejón R, Mellor-Pita S, Yebra-Bango M, Romero Y, Vargas JA. Prognostic value of neutrophils and NK cells in bronchoalveolar lavage of sarcoidosis. *Cytometry B Clin Cytom*. 2006 Nov 15;70(6):416–422.
6. Villalba, ML. Center for Drug Evaluation and Research medical review(s). Application Number 761029Orig1s000 Reference ID: 3906449 March 2016
7. Berkowitz JL, Janik JE, Stewart DM, Jaffe ES, Stetler-Stevenson M, Shih JH, et al. Safety, efficacy, and pharmacokinetics/pharmacodynamics of daclizumab (anti-CD25) in patients with adult T-cell leukemia/lymphoma. *Clin Immunol*. 2014 Dec;155(2):176–187.

8. Sahraian MA, Moghadasi AN, Owji M, Maboudi M, Kosari F, McGee JC, Minagar A. Cutaneous and pulmonary sarcoidosis following treatment of multiple sclerosis with interferon- $\beta$ -1b: a case report. *J Med Case Rep*. 2013 Dec 13;7:270.
9. Thachil J, Jadhav V, Gautam M, McKew S, Arumainathan A, Collins D, et al. The development of sarcoidosis with the use of alemtuzumab - clues to T-cell immune reconstitution. *Br J Haematol*. 2007 Aug;138(4):559-60.
10. Parisinos CA, Lees CW, Wallace WA, Satsangi J. Sarcoidosis complicating treatment with natalizumab for Crohn's disease. *Thorax*. 2011 Dec;66(12):1109-10.

## HIGHLIGHTS

- Daclizumab is a relatively new disease modifying therapy for treatment of RRMS
- A case of a 45 year old man developing sarcoidosis following daclizumab therapy
- Current understanding of sarcoid immunopathology makes its occurrence paradoxical in the setting of Daclizumab therapy
- Important to be vigilant for unexpected associations when using novel immunomodulatory therapies